

THE EISENHOWER LIBRARY



3 1151 02676 8162


STORAGE

173,264

Library



of the
Johns Hopkins University



Investigations on the Plane Quartic.

Peresa Cohen
"



*Submitted to the Board of University Studies of the
Johns Hopkins University in conformity with the require-
ments for the degree of Doctor of Philosophy.*

1918

Art. C. - 29. V. 19

	Page
1. Introduction	1
2. The Undulation	4
3. The Discriminant of the Hessian	6
4. The Discriminant of $(Sf)^4$	10
5. The Discriminant of $(tf)^6$	13
6. The Twenty-one Lines	15
7. The Invariant of Unknown Degree	17
8. Certain Other Invariants	19
9. The Eliminant of S, T, H .	25
10. The Eliminant of the Polar Cubic, Conic, and Line.	29
11. The Polar Conic of Two Points	32
12. Salmon's Conner	36
Notes	49

§ 1. Introduction.

The plane quartic $(\alpha x)^4$ is taken in the form

$$\begin{aligned} & \alpha x_0^4 \\ & + 4a_1 x_0^3 x_1 + 4a_2 x_0^3 x_2 \\ & + 6h x_0^2 x_1^2 + 12l x_0^2 x_1 x_2 + 6g x_0^2 x_2^2 \\ & + 4b_0 x_0 x_1^3 + 12m x_0 x_1^2 x_2 + 12n x_0 x_1 x_2^2 + 4c_0 x_0 x_2^3 \\ & + 4b_1 x_1^4 + 4b_2 x_1^3 x_2 + 6f x_1^2 x_2^2 + 4c_1 x_1 x_2^3 + c_2 x_2^4. \end{aligned}$$

It will be convenient first to mention briefly certain well-known forms connected with it that will be made use of in this article.

Of these several arise from the polar forms $(\alpha x)^3(\alpha y)$, $(\alpha x)^2(\alpha y)^2$, $(\alpha x)(\alpha y)^3$. Since of each of these there is an α^2 , the placing of one, two, or three conditions on the curves they represent results, respectively, in a locus for the pole, in a set of points, and in an invariant condition to be satisfied by the quartic.

Upon the polar line the only condition placeable is that it vanish identically. This, as is well known, means

that the quartic has a double point and requires the vanishing of its discriminant, an A^{27} .^{*}

The polar conic may be made to break up into two lines. The locus of poles of such degenerate conics is the Hesse in H , an A^3x^6 .^{**} To make these two lines coincide is three conditions and gives for the quartic an invariant, shown by Dr. Thomassen^{***} to be an A^{48} .

From the polar cubic one can be obtained. The locus of poles of polar cubics having a double point is the Steinerian Σ , an $A^{12}x^{12}$. The cubic has also the invariants S and T , of degrees four and six, respectively, which give rise to the covariants of the quartic $S \equiv A^4x^{14}$ [†] and $T \equiv A^6x^{16}$ ^{††}. Thus, the relations connecting the discriminant of the cubic with

* The notation $A^l x^m y^n$ is used to represent a covariant form of degree l in the coefficients of the quartic, m in x , and n in y .

** See note 1.

*** Am. Jour. Math., vol. XXXVIII, p. 249 (1916).

† See note 2.

†† See note 3.

the two invariant lines

$$\Sigma = 104 S^3 + T^2,$$

a very useful form for calculating coefficients of the Steinerian when that may be necessary. It requires that the polar cubic have a cusp is two conditions, giving rise to a set of twenty-four points with cuspidal polar cubics. The number of these points is determined by the fact that they are the common points of S and T ; therefore they are cusps of the Steinerian. There are also twenty-one points whose polar cubics have two double points and therefore break up into a line and a conic; these are the double points of the Steinerian. The line points of these cubics will hereafter be referred to simply as "the twenty-one lines." Each of these lines meets the quartic so that the four tangents at the intersection are on a pencil, which pencil is the corresponding pencil; i.e., the pencil of the cubic of which the line forms a part. If one of these lines be taken as $x_0 = 0$ and the corresponding pencil as $(1, 0, 0)$, then

$$b_0 = 24b_1 = 24b_2 = 0, \quad b_3 = 1.$$

The quartic also has certain contravariants obtained by imposing a condition on the four points in which a line cuts it. The locus of lines cutting the quartic in a self-conjugate set is $(sf)^4 \equiv s \equiv A^2 f^4$. The locus of lines cutting in harmonic pairs is $(tf)^6 \equiv t \equiv A^3 f^6$.[†]

§ 2. The Undulation.

If a line cuts the quartic in four consecutive points, the quartic is said to have an undulation. The invariant vanishing in this case is given by Salmon as an A^{60} . The undulation is evidently a line of both s and t , and it is that special case of the twenty-one lines occurring when one of the lines is on its corresponding point. This corresponding point is the undulation itself, which is therefore a double point of Σ . Suppose the undulation to be at $(0, 1, 0)$ with the tangent x_0 . Then

$$x - x_0 - f - x_1 - c.$$

[†] For the coefficients of s and t see *Am. Jour. Math.*, vol XXXIX, p. 232 (1917).

The undulation is also a point of the Hessian, for its polar conic is

$$hx_0^2 + 2b_0x_0x_1 + 2mx_0x_2 = x_0(hx_0 + 2b_0x_1 + 2mx_2),$$

a pair of lines. Let the double point of this conic be taken as $(0, 0, 1)$, which therefore becomes the corresponding point on the Steinerian. Then

$$m = 0.$$

The polar cubic of $(0, 1, 0)$ is

$$a_1x_0^3 + 3hx_0^2x_1 + 3lx_0x_1^2 + 3b_0x_0x_1^2 + 3nx_0x_2^2.$$

This is made up of x_0 and a conic; its two double points are $(0, n, b_0\sqrt{-1})$ and $(0, n, -b_0\sqrt{-1})$; they are the two points of the Hessian corresponding to the double point of the Steinerian and are harmonic to the undulation point and the Steinerian point corresponding to the undulation considered as a Hessian point. The terms in H not containing x_0 are

$$x_1^4x_2^2 - cb_0^2 + x_1^2x_2^4 - 2cb_0n + x_2^6 - cn^2 = -cx_2^2(b_0x_1^2 + nx_2^2)^2.$$

Therefore x_0 is a triple tangent to the Hessian.

§ 3. The Discriminant of the Hessian.

The Hessian and Steinerian, as is well known, are not independent curves, but are brought into one to one correspondence by the relation

$$(\alpha x)(\alpha y)^2 \alpha_i = 0, \quad i = 0, 1, 2,$$

where x is a point of the Steinerian and y a point of the Hessian. This says that the polar cubic of x has a double point y and the polar cubic of y a double point x . The joins of all such pairs x and y give rise to the Cayleyan, an $A^2 g^6$. Since x and y cannot come together unless

$$\alpha x^3 \alpha_i = 0, \quad i = 0, 1, 2,$$

which is the condition that the quartic have a double point, there is a very convenient reference scheme for handling these three curves in any other case. Let $(0, 1, 0)$ be a point of the Hessian, $(0, 0, 1)$ the corresponding point on the Steinerian, so that x_0 is a line of the Cayleyan. Then

$$\alpha x^3 \alpha_i = 0, \quad i = 0, 1, 2,$$

$$g_0 = \alpha x^3 \alpha_i = 0.$$

Now let us see under what conditions the Steinerian requires a double point. It is known to have one when the quartic does. To discover other cases let us use the above reference scheme. Then

$$H = x_1^5 x_0 \cdot 2n(bh - b_0^2) + x_1^5 x_2 \cdot 2c_1(bh - b_0^2) \\ + \text{lower terms in } x_1.$$

Therefore $(0, 1, 0)$ is a double point if

$$bh - b_0^2 = 0.$$

But this says that the polar conic of $(0, 1, 0)$, which is

$$hx_0^2 + 2b_0 x_0 x_1 + bx_1^2,$$

shall be a line counted twice, and this means the vanishing of an A^4 . Since this line is already on $(0, 0, 1)$, let it be taken as x_1 , so that

$$b_0 = b = 0.$$

Then the polar cubic of any point $(K, 0, 1)$ on x_1 , namely,

$$(aK + a_2)x_0^3 + 3(a_1K + l)x_0^2x_1 + 3(a_2K + g)x_0^2x_2 + 6(lK + n)x_0x_1x_2 \\ + 3(gK + c_0)x_0x_2^2 + 3(nK + c_1)x_1x_2^2 + (c_0K + c)x_2^3,$$

has a double point at $(0, 1, 0)$; therefore the Steinerian must contain the line x_1 . Two of these polar cubics have cusps, the

poles of which may be made $(1, 0, 0)$ and $(0, 0, 1)$ by requiring that

$$l = n = 0.$$

The cuspidal tangent of each cubic is on the pole of the other, and the pair of them make up the tangents to the Hessian at its double point. Then

$$S = -a_1^2 c_1^2 x_0^2 x_2^2 - \dots - x_1 \left[x_0^3 (-b a_1 a_2 c_0 + b a_1 g^2) + \dots + x_2^3 (-b a_2 c_0 c_1 + b c_1 g^2) \right] + \text{higher terms in } x_1,$$

and

$$T = 8 a_1^3 c_1^3 x_0^3 x_2^3 + x_1 \left[x_0^5 \cdot 4 b a_1^3 c_0^2 + \dots + x_2^5 \cdot 4 b a_2^2 c_1^3 \right] + \text{higher terms in } x_1.$$

Therefore S touches x_1 , and T has x_1 as a flex line at $(1, 0, 0)$ and $(0, 0, 1)$. Also

$$\Sigma = x_1 \left[x_0^8 x_2^3 \cdot 64 b a_1^6 c_0^2 c_1^3 + \dots + x_0^3 x_2^8 \cdot 64 b a_1^3 a_2^2 c_1^6 + x_1 \left\{ x_0^{10} \cdot 16 b^2 a_1^6 c_0^4 + \dots + x_2^{10} \cdot 16 b^2 a_2^4 c_1^6 \right\} + \text{higher terms in } x_1 \right],$$

showing that x_1 divides out only once, but that $(1, 0, 0)$ and $(0, 0, 1)$

are more than ordinary singularities.

The Hessian also has a double point when

$$c_1 = 2c_0 = 0.$$

The symmetry shows that $(0, 0, 1)$ is also a double point, as can be verified from the coefficients. Then $(0, 1, 0)$ as a Hessian point has $(0, 0, 1)$ as the corresponding Steinerian point, and vice versa. The polar conics of $(0, 1, 0)$ and $(0, 0, 1)$ are, respectively,

$$ax_0^2 + 2b_0x_0x_1 + cx_1^2$$

$$\text{and } gx_0^2 + 2c_0x_0x_2 + cx_2^2.$$

If the harmonic conjugate of x_0 as to the first pair of lines is taken as x_1 , and that as to the second as x_2 , then

$$b_0 = c_0 = 0.$$

This shows that x_0 is one of the twenty-one lines. To find the degree of the invariant consisting in this case is one of the objects of this investigation.

With the above reference scheme

$$J = x_1x_2^2 - b_0x_1^2$$

$$+ x_2^2 [x_0^2 (cx_1^2 - 2bx_0x_1) + x_0x_1 (-bca_1 + x_1^2 (-bca_1 - 5cb^2))] \\ + 10ab_0x_0x_1x_2,$$

$$T = x_1 x_2^5 \cdot 2 b c^2 l^3$$

$$+ x_2^4 \left[x_0^2 \cdot -3 c^2 h^2 l^2 + 2 x_0 x_1 (-3 b c^2 a_2 h l + 6 b c^2 a_1 l^2) \right.$$

$$\left. + x_1^2 (b^2 c^2 a_2^2 + 4 b^2 c g^3 + 12 b c^2 h l^2) \right] + \text{lower terms in } x_2.$$

Therefore these two curves touch at $(0, 0, 1)$ along the line x_1 , and similarly at $(0, 1, 0)$ along the line x_2 .

$$Z = x_1^2 x_2^{10} + b^2 c^4 l^6$$

$$+ x_2^9 \left[x_0^2 x_1 - 12 b c^4 l^2 l^5 + x_0 x_1^2 \cdot b c l^3 (-3 b c^2 a_2 h l + 6 b c^2 a_1 l^2) \right.$$

$$\left. + x_1^3 \cdot 4 b c^2 l^3 (b^2 c^2 a_2^2 - 12 b^2 c g^3 + 12 b c^2 h l^2) \right] + \text{lower terms in } x_2.$$

Therefore Z has a singularity at $(0, 0, 1)$, and symmetrically at $(0, 1, 0)$, which is something more than merely a cusp. It may be a tac node.

These three cases make up the totality of ways in which the Hessians may acquire a double point. Therefore its discriminant, A^{225} , must be made up of the three varieties attached to the ...

§ 4. The Discriminant of $(5f)^4$

The contravariant $(5f)^4$ is on the line x_0 if

$$b_0 = f - c - c.$$

The point of contact is given by

$$3c_1 m J_1 + (-b c_0 + b_0 c_1) J_2 = 0.$$

Since $(0, 0, 1)$ is taken as any one of the points in which x_0 cuts the quartic, this point of contact will be on $(\alpha x)^4$ if

$$m = 0. \quad +$$

But this says that $(0, 0, 1)$ is a point of the Steinerian. Therefore the forty-eight intersection of $(\alpha x)^4$ and $(s f)^4$, which considered as a point curve is of order twelve, are the same as those of $(\alpha x)^4$ and Σ . This can be substantiated by finding the

point equation of $(s f)^4$. The line equation of $(\alpha x)^4$ is $s^3 - 27 t^2$.

But s formed for $(s f)^4$ is $-12 S + A^3 (\alpha x)^4$, where A^3 is the invariant A given by Salmon, and t formed for $(s f)^4$ is

$T + (\alpha x)^4 \cdot (t \alpha')^3 (t \alpha'')^3 (\alpha' x) (\alpha'' x)$. Therefore the line equation of $(s f)^4$ is

$$\begin{aligned} [-12 S + A^3 (\alpha x)^4]^3 - 27 [T + (\alpha x)^4 \cdot (t \alpha')^3 (t \alpha'')^3 (\alpha' x) (\alpha'' x)]^2 \\ = -27 \Sigma + (\alpha x)^4 \cdot A^3 x^8. \end{aligned}$$

x_0 will become a double line of S only if

$$b_0 - c_0 = m = 0,$$

⁺ If $c_1 = 0$, the quartic would have an undulation.

all other conditions than this placing more than one restriction on the quartic. The points in which x_0 meets the quartic are $(0, 0, 1)$, $(0, K, 1)$, $(0, \omega K, 1)$, $(0, \omega^2 K, 1)$, where $K^3 = -\frac{4c_1}{b}$ and $\omega^3 = 1$. The tangents to the quartic at these four points are, respectively,

$$c_1 x_1 = 0,$$

$$3\pi K x_0 - 3c_1 x_1 + 3c_2 K x_2 = 0,$$

$$3\pi \omega K x_0 - 3c_1 x_1 + 3c_2 \omega K x_2 = 0,$$

$$3\pi \omega^2 K x_0 - 3c_1 x_1 + 3c_2 \omega^2 K x_2 = 0,$$

and they evidently have the common point $(c_1, 0, -\pi)$. Let

$$x = 0,$$

so that this common point becomes $(1, 0, 0)$. Since among the coefficients equated to zero are $\alpha_2, \alpha_3, \beta$, then $(0, 1, 0)$ is a point of the Hessian with $(0, 0, 1)$ as its corresponding Steinerian point. Since the tangent to the Steinerian at any point is the polar line as to the quartic of the corresponding Steinerian point, the tangent to the Steinerian at $(0, 1, 0)$ is x_1 , which is the tangent to x_1^4 at that same point. Thus since no distinction was made among the four points

in which x_0 cuts the quartic by assigning the particular coordinates $(0, 0, 1)$ to one of them, when one of the twenty-one lines is a line of $(\mathcal{S}\mathcal{F})^4$, the quartic and Steinerian touch four times on this line, the four tangents going thru the corresponding point. Also, since the tangent to the Hessian at $(0, 1, 0)$ is x_2 , the tangents at the four Hessian points corresponding to these Steinerian points are also on that corresponding point.

x_0 will also be a line of $(\mathcal{S}\mathcal{F})^4$ if

$$b = b_2 = f = 0;$$

i.e., if x_0 is a stationary line of the quartic. But this leads to nothing new. Therefore $(\mathcal{S}\mathcal{F})^4$ has a double line only when one of the twenty-one lines is on $(\mathcal{S}\mathcal{F})^4$, and its discriminant, an A^{54} , expresses this condition.

§ 5. The Discriminant of $(t\mathcal{F})^6$.

The contravariant $(t\mathcal{F})^6$ is on the line x_0 if

$$b = b_2 = f = 0.$$

Then

$$(t\mathcal{F})^6 = 2\mathcal{F}_0^5\mathcal{F}_1 \cdot 2b_0c_1^2 + \text{lower powers in } \mathcal{F}_0.$$

x_0 becomes a double line either if $b_0 = 0$, which says that the quartic has a double point at $(0, 1, 0)$ with x_0 as one of the tangents there, or if $c_1 = 0$, which says that the quartic has an inflection at $(0, 1, 0)$.

$(t\mathcal{F})^6$ is also on x_0 when

$$b_2 = f = c_1 = 0.$$

Then

$$(t\mathcal{F})^6 = 2\mathcal{F}_0^5\mathcal{F}_1 \cdot -bcn + 2\mathcal{F}_0^5\mathcal{F}_2 \cdot -bcm + \text{lower powers in } \mathcal{F}_0.$$

For x_0 to be a double line requires either $b=0$ or $c=0$, which repeats the undulation condition, or $m=n=0$, which is the condition on the quartic under which the Hessian has two double points and for which the invariant of unknown degree vanishes. Then one of the twenty one lines is a line of $(t\mathcal{F})^6$.

$(t\mathcal{F})^6$ also has x_0 as a line when

$$b = f = c = 0,$$

but this leads to nothing new. Therefore the discriminant

of $(t\mathfrak{f})^6$, an A^{225} , must be made up of the discriminant of the quartic, an A^{27} , the undulation condition, an A^{60} , and the unknown invariant.

§ 6. The Twenty-one Lines.

The twenty-one lines* are given by an $A^i x^{21}$, where i is as yet unknown. That it can be determined is due to the fact that these lines are part of the common lines of two curves, all of whose common lines are known. They are, the Cayleyan, an $A^{12} \mathfrak{f}^{18}$, and an $A^{15} \mathfrak{f}^{24}$, which is the locus of lines cutting the quartic so that three of the tangents at the intersections are on a point.** The common lines of these curves are: *** 1) the twenty-one lines counted sixteen times

* A curve in three lines is

$$2(s\mathfrak{f})^3(s\alpha)(s'\mathfrak{f})(s'\alpha)(s''\mathfrak{f})(s''\alpha)(s'''\mathfrak{f})(s'''\alpha) - 3(s\mathfrak{f})^4(s'\mathfrak{f})^3(s'\alpha)(s'\mathfrak{f})(s'\alpha)(s'''\mathfrak{f})(s'''\alpha)^2 \\ + \frac{1}{3}(s\mathfrak{f})^4(s'\mathfrak{f})^4(s''\mathfrak{f})^4(s'''\mathfrak{f})^4.$$

** Am. Jour. Math., vol. XXXIX, p. 227 (1917).

*** Let $t_2 = m = \mathfrak{f} = 0$, so that x_0 is a line of the Cayleyan. Let

since they are quadruple lines of both curves; 2) the twenty-four stationary lines, given by an $A^{24}x^{24}$, counted twice because they are double lines of $A^{15}f^{24}$; 3) the forty-eight lines of sf^{12} at its intersection with the quartic. Since the polar points of these lines are on $(\alpha x)^4$, they are lines of the curve $(sf)^3(s\alpha)(s'f)^3(s'\alpha)(s''f)^3(s''\alpha)(s'''f)^3(s'''\alpha)$, an A^9f^{12} , and are obtained by the elimination of f from A^9f^{12} , $sf^{12} = A^2f^4$, and (αx) as in $A^{60}x^{48}$. The eliminant of $A^{12}f^{18}$, $A^{15}f^{24}$, and (αx) is an $A^{558}x^{432}$, which must be made up of the same old lines together with

$n=0$, so that x_2 is tangent to the Hessian at $(1,0,0)$. The tangent to the discriminant cannot be chosen as the other inflexion line, because the stationary lines are known to be among the common lines of the two curves and the Steinerian is tangent to them. Instead let $(1,0,0)$ be determined by the intersection of x_2 with the polar line of $(0,0,1)$. Then $c_0=0$. The coefficient of f_0^{24} in $A^{15}f^{24}$ becomes $\frac{12}{3}b^2c^2b^4c^8$. If $b=0$, x_0 is a stationary line of the quartic. If $c=0$, the quartic and Σ meet at $(0,0,1)$ and x_0 is a line of sf^{12} . If $b_0=0$, x_0 is one of the twenty-four lines. If $c_0=0$, an involution condition is imposed on the quartic.

such invariants as express the fact that they become indeterminate. The only such invariant is the discriminant of the quartic, for if the quartic has a double point it divides out at least once from each of these curves. Therefore

$$A^{558} x^{432} = (A^6 x^{21})^{16} \cdot (A^{14} x^{21})^2 \cdot A^{10} x^{118} \cdot (A^{27})^k$$

$$\therefore 558 = 16i + 108 + 27k,$$

$$16i + 27k = 450.$$

This equation shows that i must be a multiple of nine and can be satisfied only when $i = 18$. Therefore the twenty-one lines are given by an $A^{18} x^{21}$.

§ 7. The Invariant of Unknown Degree.

The degree of the form giving the twenty-one lines being known, we can force one of them to lie on either $(sf)^{10}$ or $(tf)^6$. The condition so obtained will contain the undulation condition, however, for the tangent at an undulation is a special case of one of the twenty-one lines and is a line of both s and t . Suppose the $A^{18} x^{21}$ factored into its component

lines, the coordinates of each of these substituted in $(t\mathcal{F})^4$, the twenty-two resulting terms multiplied together, and the coefficients of the $A''^8 x^{21}$ substituted for symmetric functions of these coordinates. The result is

$$A^{18 \cdot 4 + 21 \cdot 21} = A^{114} = A^{60} \cdot A^{54}.$$

This leads to nothing new. But the same process applied to $(t\mathcal{F})^6$ gives the degree of the required invariant. For we obtain $A^{18 \cdot 6 + 21 \cdot 3} = A^{171}$ from which the undulation condition must divide out at least once. Therefore the invariant in question may be either of degree 111 or of degree 51. The first of these will not fit into the discriminant of the Hessian, but the second would give

$$A^{225} = A^{27} \cdot (A^{48})^2 \cdot (A^{51})^2,$$

while for the discriminant of $(t\mathcal{F})^6$ we should have

$$A^{225} = (A^{21})^2 \cdot (A^{60})^2 \cdot A^{51}.$$

Therefore we have an invariant of degree 51, the vanishing of which expresses the fact that the Hessian has 110 double points, that $(t\mathcal{F})^6$ has a double line, and that one of the twenty-one lines is a line of $(t\mathcal{F})^6$. A further meaning will be seen later.

§ 8. Certain Other Invariants.

The degree of the form giving the twenty-one lines being known, that for the twenty-one corresponding points can be calculated from the fact that each point is the polar point of the corresponding line as to both SY^4 and TJ^6 . Suppose, as before, the $A^{18}x^{21}$ factored into its component lines, the coordinates substituted for η in $(SJ)(S_1J)^3$, and the symmetric functions of these coordinates in the product of the resulting terms replaced by the coefficients of $A^{18}x^{21}$. The result gives the twenty-one points together with such conditions as express that the polar point becomes indeterminate (i.e., that one of the lines be a double line of S). Then

$$A^{18 \cdot 3 + 2 \cdot 21} g^{21} = A^{54} \cdot A^{42} g^{21}.$$

Similarly from $(tJ)^6$ we obtain

$$A^{18 \cdot 5 + 3 \cdot 21} g^{21} = A^{60} \cdot A^{51} \cdot A^{42} g^{21}.$$

Therefore the twenty-one points are given by $A^{42} g^{21}$. *

* This degree fits in with the form $A^{36} g^{18}$ for the line equation of Σ . For the cusps are obtained from the discriminant of S, T, Jx as an $A^{48} g^{24}$. Then from the Plücker formula

$$A^{12 \cdot 22} g^{12 \cdot 11} = A^{264} g^{132} = (A^{42} g^{21})^2 \cdot (A^{48} g^{24})^3 \cdot A^{36} g^{18}.$$

one of them is $\text{con}(ax)^4$ (and therefore so is its reciprocal cubic). Then, as before, considering $A'^{12}y^{21}$ broken up into its component points, the coördinates substituted in $(ax)^4$, and their reciprocal functions replaced in the product of the resulting terms, we obtain an $A'^{42 \cdot 4 + 21} = A'^{89}$. Out of this the condition for an undulation, when the point is on the line part of its polar cubic, must divide out at least once. If it is no more than once, we have left an A'^{29} to express that the point is on the conic part of its polar cubic. To substantiate this degree we must find some case into which the undulation condition does not enter. We had before that the point is $(1, 0, 0)$ and its corresponding line x_0 if

$$b_0 = m = n - c_0 = c.$$

To put the point on the quadratic requires that

$$x = 0.$$

The undulation case is excluded because our choice of coördinates for the point and line implies that they are not incident. We can still choose $(0, 1, 1)$ and $(0, 0, 1)$ in x_0 .

Let us take them as the double points of the polar cubic of $(1, 0, 0)$, so that

$$q = r = 0.$$

Then, in $(sf)^4$ the coefficients of $J_1^4, J_2^4, J_1^3 J_2, J_1 J_2^3$ vanish, showing that x_1 and x_2 are lines of s with contact $(1, 0, 0)$, so that $(1, 0, 0)$ is a double point of $(sf)^4$.* It is known that the double lines of $(sx)^4$ are given by $A^{16} x^{28}$; therefore the double points of $(sf)^4$ are given by an $A^{32} f^{28}$. One of them can lie on $(sx)^4$ only in our present case or when $(sx)^4$ has a double point, for then $(sf)^4$ has that same double point with the same tangents. Then by the method of considering the $A^{32} f^{28}$ factored into its component parts, we have as the condition that one of these points be on $(sx)^4$

$$A^{32 \cdot 4 + 28} = A^{156} = A^{129} \cdot A^{27}.$$

This establishes the degree of the A^{129} .

The totality of polar cubics of the twenty-one points given by the $A^{42} f^{21}$ is given by an $A^{63} x^{63}$. Suppose this

* An examination of the coefficients of $(sf)^4$ shows that an undulation is not a double point of it.

factored into its component cubics $(p_i x)^3$, and then each of these cubics factored into its component line and conic $(y_i x) \cdot (\delta_i x)^2$. Then the condition that the line touch the conic is of second degree in both y_i and δ_i , therefore of second degree in p_i . Therefore the invariant expressing the condition that there be a polar cubic made up of a conic and a line touching it is an $A^{6 \cdot 3 \cdot 2} = A^{12 \cdot 6}$. This result can be substantiated in other ways. If, as usual, we take the line as x_0 and the corresponding point as $(0, 0, 0)$, so that

$$b_0 = m = n = c_0 = 0,$$

then the conic will touch x_0 at $(0, 1, 0)$ if

$$gh - l^2 = 0.$$

Then S goes thru $(1, 0, 0)$ while T has a double point there, so that Z has ^a~~two~~ double points and two cusps coming together and acquires a triple point, since x_0^9 is the highest power of x_0 recurring in its equation. Since two of the intersections of S and T have come together, the $A^{12 \cdot 6}$ must be a factor of the tact-invariant of these two curves. We have seen (p. 8, 10) that in case of the vanishing of either

the A^{48} or the A^{51} that S and T touch at two distinct points, and these three cases seem to be the only ones in which two intersections of these curves come together. Their tact-invariant is found from the formula given by Salmon to be of degree $4 \cdot 6(6+8-3) + 6 \cdot 4(4+12-3) = 576$. Then

$$A^{576} = (A^{48})^2 \cdot (A^{51})^2 \cdot (A^{126})^3.$$

Also inspection of their coefficients shows that both H and S touch x_0 at $(0, 1, 0)$. As will be seen later, two intersections of S and H can come together only when two of S and T do. When the A^{51} vanishes, H has two double points on S ; when the A^{48} vanishes, H has a double point which is also a double point of S . The tact-invariant of S and H is of degree $4 \cdot 6(6+8-3) + 3 \cdot 4(4+12-3) = 420$.

$$A^{420} = (A^{48})^4 \cdot (A^{51})^2 \cdot A^{126}.$$

The twenty-one polar cubics being given by an $A^{63}x^{63}$, while the twenty-one line parts are given by an $A^{18}x^{21}$, the twenty-one conic parts are given by an $A^{45}x^{42}$. Since the discriminant of a conic is of third degree in its coefficients, the condition that one of these conics break up

is an A^{45} . But then the polar cubic consists of three lines, any two of which may be considered as the conic part, so that this degree is probably divided by three. Therefore the condition that there be a polar cubic made up of three lines is an A^{45} . Then three of the double points of the Steinerian have come together, giving it a triple point.*

* Professor Nozley has pointed out to me that the degrees of these last two invariants can be substantiated by the work of Caporali (*Memorie di Geometria*, p. 111) on a web of plane curves. If i curves of a web are of a particular kind, then the invariant expressing that there be a curve of that kind in a web is of degree i in the coefficients of the three curves upon which the web is built up. From his results the invariants expressing that in a web of cubics there be a cubic with three double points or a cubic with two coincident double points, are of degree 10 and 12, respectively, in the coefficients of the three cubics. But in a polar web each curve is linear in the coefficients of the quartic. Therefore the invariants are of degree 40 and 120 in the coefficients of the quartic.

§ 9 The Eliminant of S, T, H

Under the reference scheme there

$$b_2 = f - c - o,$$

the coefficient of the highest term in x_2 in both S and T is a power of $c, b+n^2$. Then both coefficients vanish if $c, = n = 0$, but this, as we have seen, means the vanishing of an A^5 . Then for the general quartic we obtain $(0, 0, 1)$ as one of the common points of S and T if we make

$$b = n = 0.$$

But then the coefficient of x_1^4 vanishes in S . Therefore the twenty-four roots of the determinantal correspond three twenty-four points of the Hessian which make up its intersections with S .

It is known that the polar cubic of a point on S can be reduced to the sum of the cubes of three linear factors; its Hessian is the product of these factors and is one of three four triangles whose sides pass through all the flexes of the cubic. If the cubic has a double point ^{proper} there is only one such triangle made up

of the tangents at the double point and a line thru the three remaining flexes (flex line); in case of a cusp even this degenerates to the cusp tangent counted twice and the line joining the cusp to the sole remaining flex.

Here the polar cubic of $(0, 0, 1)$ is

$$a_2 x_0^3 + 3g x_0^2 x_2 + 3c_0 x_0 x_2^2 + 3c_1 x_1^2 x_2 + c_2 x_2^3,$$

and its Hessian is

$$c_1^2 x_2^2 (a_2 x_0 + g x_2)$$

But x_2 is the tangent to H at $(0, 1, 0)$ and $a_2 x_0 + g x_2$ is the tangent to S at the same point. Therefore the polar cubic of a cusp of the Steinerian has as its cusp tangent and its flex line the tangents to H and S , respectively, at the corresponding point.

The cubic can be thrown into the form

$$\frac{1}{a_2^2} \left[(a_2 x_0 + g x_2)^3 + x_2^2 \left\{ 3a_2 (a_2 c_0 - g^2) x_0 + 3a_2^2 c_1 x_1 + (ca_2^2 - g^3) x_2 \right\} \right]$$

which also shows up the tangent at the flex as the linear form in brackets

The polar cubic of $(0, 1, 0)$ is

$$a_1 x_0^3 + 3h x_0^2 x_1 + 3b_0 x_0 x_1^2 + b x_1^3 + c_1 x_2^3,$$

and its Hessian is

$$c_1 x_2 \left[x_0^2 (a_1 b_0 - h^2) + x_0 x_1 (b a_1 - b_0 h) + x_1^2 (b h - b_0^2) \right].$$

Then the tangent to H at $(0, 1, 0)$ passes thru three flexes of the cubic, the three flex tangents pass thru the corresponding Steinerian point and the binary Hessian of these three tangents picks up the six remaining flexes.

The coefficient of x_2^6 in H is $-c_1^2 g$. Since we cannot have $c_1 = 0$ without bringing on the vanishing of the A^{51} and worse, let us make

$$g = 0.$$

Then we have made S, T, H have a common point at $(0, 0, 1)$.

The polar cubic of $(0, 0, 1)$ becomes

$$a_2 x_0^3 + x_2^2 (3 c_0 x_0 + 3 c_1 x_1 + c_2 x_2).$$

If we let the intersection of the stationary tangent and flex line be $(1, 0, 0)$, then

$$c_0 = 0.$$

But then, since we have $g = n = c_0 = 0$, the Steinerian point corresponding to $(0, 0, 1)$ as a Hessian point is $(1, 0, 0)$, which also becomes a cusp on the Steinerian. So, to sum up, we have: 1) the cusp e_0 on the Steinerian corresponds to e_2 on

the Hessian; 2) the cusp e_2 on the Steinerian corresponds to e_1 on the Hessian; 3) the tangent to the Hessian at e_1 goes thru e_0 and at e_2 goes thru e_1 ; 4) the polar cubic of e_2 has the tangent to the Hessian at e_1 as ~~an~~ tangent, the line $x_0 (= e, e_2)$ as flex line, and a stationary tangent going thru e_0 ; 5) the polar cubic of e_2 has an analogous set of lines; 6) S is tangent to x_0 at e , and to x_1 at e_2 .

The real underlying significance of this state of affairs is not ~~the same~~; it is the degree of the invariant before setting it, but it now passes at. The discriminant of S, T, H is an A^{360} . The only other condition under which these three curves have a common point is when the A^{37} vanishes, but then they have four common points, for H has two double points on the other curves. Then, presumably, the A^{37} divides four times out of the discriminant, leaving an A^{356} . The essential difference between the two cases seems to be this: To every ST point (intersection of S and T), there is attached an SH point, with which it cannot coincide without some special condition on the quartic. I shall for S, H



It is known a common point is an ST point, and it coincides with an SH point. If then the corresponding SH and ST points also coincide, the H^2 coincides. There there is no further coincidence as in the other case.

§10 The Eliminant of the Polar Cubic, Conic, and Line.

Suppose we ask that the polar cubic, conic, and line of $(0, 1, 0)$, which is not on the quartic, have a common point $(0, 0, 1)$.

Then

$$b_2 = f = c = 0.$$

The symmetry of this condition shows that the relation ~~set~~ between $(0, 1, 0)$ and $(0, 0, 1)$ is a mutual one. Also, x_0 is a line of H^2 . Therefore we might define $(H^2)^6$ as the locus of the joins of pairs of points so related that the polar curves of each pass thru the other, the pair of points being the common harmonic pair of the two harmonic pairs of points in which their join cuts the quartic.

In general the eliminant of the polar curves of x , which

are an Ax^3y , an Ax^2y^2 , and an Axy^3 , is an $A''x^{26}$. Out of this x^{12} divides twice, leaving an $A^9x'^9$. Before extracting the meaning of this latter curve, let us examine the eliminant of a cubic Ay^3 , its polar cubic Axy^2 , and its polar line Ax^2y . It is an $A''x^{14}$, from which the cubic divides out twice. The remainder, an $A^9x'^9$, gives the nine stationary lines. Therefore the $A^9x'^9$ is for the quartic the locus of points lying on the stationary lines of their polar cubics, also the locus of flexes of polar cubics the tangent at which lies on the cubic.

Salmon gives as the equation of the nine stationary lines of the cubic

$$5S U^2 H - H^3 - U O,$$

where S is the invariant of degree four, U is the cubic itself, H is its Hessian, and O is a certain A^8x^6 . For a polar cubic of the quartic the first three pass over readily enough into the covariant S , the quartic itself, and its Hessian, but O is not so easily transferred. So we will substitute for it another A^8x^6 , expressible in terms of S , U , H , and O , which gives the locus of points whose polar conics as to the cubic

are on their polar lines as to the Hessian. Then it is easily shown that

$$A^9 x^{18} = S U^2 H^3 + 9 U O',$$

where S is the covariant S of the quartic, U the quartic, H its Hessian, and O' the locus of those points whose polar conics as to the quartic are on their polar lines as to the Hessian.

The form of the $A^9 x^{18}$ shows that all of its intersections with U^4 are used up at the flexes of the quartic. These twenty-four points are also part of its intersections with H , but there are eighty-four more, lying on O' . Now we saw that $(0, 1, 0)$ is a point of the $A^9 x^{18}$ if

$$b_2 = f = c = 0.$$

To make it a point of H also requires that

$$m = 0. \quad *$$

But there are only forty-two such points, for they are the points where H, U^4 , and the Cayleyan all touch. ** Then the intersections of H and O' must come together by two's, and,

* Or, symmetrically, $(0, 0, 1)$ is a point of H if $n = 0$.

** Proc. Nat. Ac. Sci., vol. 3, p. 449 (1917)

indeed, it is readily shown that the two curves touch at their common points.

It is also easily shown that the forty-two inflexions of the Steinerian are at the points corresponding to these Hessian points.

§ 11. The Polar Conic of Two Points.

The eliminant of the polar curves of the quartic connects up with the polar conic of two points in the following way. The polar conic of x and y as to the quartic is $(ax)(ay)(x^2y^2)$. Any point on the line x, y may be represented parametrically as $x + \lambda y$. This line meets the conic in two points whose parameters λ are given by

$$(ax)(ay)(x + \lambda y)^2 = (ax)^3(ay) + 2\lambda(ax)^2(dy)^2 + \lambda^2(ax)(dy)^3 = 0.$$

If we ask that these points become indeterminate, we ask that the coefficients of λ all vanish. Then x and y are a pair of points such that the polar curves of each are on the other, so that we can define the $A^9x'^2$ of the last section as the locus

of pairs of points whose polar conic is not only degenerate but contains their join, and $l f^6$ as the locus of this join. If f is now taken as $(0, 1, 0)$ and $(0, 0, 1)$, then, as we have seen,

$$b_2 = f = c_1 = 0.$$

The polar conic of the two points is

$$lx_0^2 + 2mx_0x_1 + 2nx_0x_2 = x_0(lx_0 + 2mx_1 + 2nx_2).$$

To make the double point at one of the points requires that

$$m = 0 \quad (\text{or } n = 0).$$

But this makes $(0, 1, 0)$ one of the forty-two points on H where it touches $(cf)^6$.

In general for a given f we have a whole locus of points x , the polar locus of f to be exact, such that the polar conic of x and f is a pair of lines. We have seen that for one of these lines to be xy , x and f must lie on an $A^9x'^2$. If we ask instead that the double point be x , then x is a point of the Steinerian and f a point of the distribution. For the polar conic of $(0, 1, 0)$ and $(0, 0, 1)$ is

$$lx_0^2 + 2mx_0x_1 + 2nx_0x_2 + b_2x_1^2 + 2fx_1x_2 + c_1x_2^2,$$

and this becomes two lines on $(0, 1, 0)$ if

$$x_2 - f = 0,$$

being, in fact, the tangents at the double point of the polar cubic of the Steinerian point. To see both that x, y be one of the lines and that the double point be at x facts, it, as we have seen, the forty-two inflexions of the Steinerian and the forty-two points where H touches the tangents and $(6f)^4$.

Making a fresh start, let us ~~ask~~ that the polar conic of $(0, 1, 0)$ and $(0, 0, 1)$ be the square of a line. If we take $(1, 0, 0)$ as a point of this line, then

$$x - y - z = 0,$$

$$x_2 c_1 - f^2 = 0.$$

But then we find that the coefficients of x_1^4 and x_2^4 in S vanish. Therefore the pair of points whose polar conic is a repeated line lie on S . But we already have a correspondence of the points of S , for the polar Hessian of a point of S is made up of three lines, the intersections of which are on S . With the above reference scheme

$$S_{4,10} = (x_1 c_0 - h^2)(-x_2 c_0 + f^2).$$

$(1, 0, 0)$ was any point of the repeated line. Let us take it as one of the intersections with S by making

$$a, b_0 - h^2 = 0.$$

Then the polar-Hessian of $(0, 1, 0)$ becomes

$$x_1 \left[x_0 x_1 (b a_1 f - a, b_2^2 - b_0 f h) + x_0 x_2 (b a_1 c_1 - a, b_2 f - b_0 c_1 h) \right. \\ \left. + x_1^2 (b f h - b_0^2 f - b_2^2 h) + x_1 x_2 (b c_1 h - b_0^2 c_1 - b_2 f h) \right].$$

The double points are $(1, 0, 0)$, $(0, 0, 1)$ and

$$y \equiv (b c_1 h - b_0^2 c_1 - b_2 f h, -b a_1 c_1 + a, b_2 f + b_0 c_1 h, b a_1 f - a, b_2^2 - b_0 f h).$$

Then the polar conic of $(0, 1, 0)$ and $(1, 0, 0)$ becomes $a, x_0^2 + 2 h x_0 x_1 + b_0 x_1^2$, that of $(0, 1, 0)$ and $(0, 0, 1)$ is $b_2 x_1^2 + 2 f x_1 x_2 + c_1 x_2^2$, and that of $(0, 1, 0)$ and y is x_1^2 . Therefore the polar conic of a point on S and any vertex of its polar-Hessian triangle is the square of the opposite side. *

* We might expect this sort of three-to-one correspondence from the following considerations: The line equation of any yz^2 is an $A^2 x^2 y^2 z^2$. If the conic is the square of a line, the line equation vanishes identically. Equating each coefficient to zero, we have six equations from which we can eliminate the six quantities $y_0^2, y_1^2, y_2^2, y_1 y_2, y_2 y_0, y_0 y_1$, giving an $A'^2 x'^2$. This must be S^3 .

If now we also ask that the repeated line be one $(0, 1, 0)$ we have

$$b_2 = f = 0.$$

Then $(0, 1, 1)$ is one of the twenty-four intersections of S and H . Therefore the polar conic of an SH point and its corresponding ST point is the square of a line on the SH point & the tangent to H there.

If, finally, we ask that the polar conic of $(0, 1, 1)$ and $(0, 0, 1)$ be x_2^2 , we have

$$a_1 = a_2 = b_2 = f = c_1 = 0.$$

But then the $A^{(5)}$ vanishes. Therefore we have a new definition for the $A^{(5)}$; it is the invariant expressing the condition that there be a frame of points such that their polar conic as to the quartic is the square of the line joining them.*

§12. Salmon's Connex.

Salmon has shown that with any plane curve $(2s)^n$ there

* See note 4.

is associated an $A^3 x^{2(n-2)} y^{n-2}$, which when x is a point on the curve picks out the remaining intersections of the tangent at x with the curve. This connex is expressible in terms of polars of the Hessian of the curve itself and the Hessians of the polar curves of x as to $(ay)^n$. In case of the quartic the connex is an

$$A^3 x^4 y^2 \equiv 15(hx)^4(hy)^2 - 9(h, x)(h, y)^2,$$

where $(hy)^4$ is the Hessian of the quartic and $(h, y)^3$ is the Hessian of the polar cubic of x . Explicitly

$$\begin{aligned} A^3 x^4 y^2 = \sum^3 \left[x_0^4 \{ y_0^2 \cdot 6(a_0 h - a_1^2 - a_1^2 g - a_2^2 h + 2a_1 a_2 l) \right. \\ + y_0 y_1 \cdot 4(ab_0 g + ahn - 2alm - a_2^2 b_0 - a_1^2 n \\ + 2a_1 a_2 m - a_1 gh + a_1 l^2) \\ + y_0 y_2 \cdot 4(ac_0 h + agm - 2aln - a_1^2 c_0 + 2a_1 a_2 n \\ - a_2^2 m - a_2 gh + a_2 l^2) \\ + y_1^2 (abg - ba_2^2 + ab_0 n - 2ab_2 l + afh - am^2 \\ + 2a_1 a_2 b_2 - a_1^2 f - a_1 b_0 g - a_1 hn + 2a_1 lm) \\ + y_1 y_2 (ab_0 c_0 + 2ab_2 g + 2ac_1 h - 4afl - amn - 2a_1^2 c_1 \\ + 2a_2^2 b_2 + 4a_1 a_2 f - a_1 c_0 h - a_2 b_0 g \\ \left. - a_1 gm - a_2 hn + 2a_1 l n + 2a_2 l m) \right] \end{aligned}$$

$$\begin{aligned}
& + y_2^2 (a_1^2 h - c a_1^2 + a c_0 n - 2 a c_1 l + a f g - a n^2 + 2 a_1 a_2 c_1 \\
& \quad - a_2^2 f - a_2 c_0 h - a_2 g m + 2 a_2 l n) \} \\
& + x_0^3 x_1 (f_0^2 \cdot 8 (a b_0 g + a h n - 2 a b_1 n - a_2^2 b_0 - a_1^2 n + 2 a_1 a_2 n \\
& \quad - a_1 g h + a_1 l^2)) \\
& + y_0 y_1 \cdot 2 (a b g - b a_2^2 + 7 a b_0 n - 2 a b_2 l + a f h - 7 a m^2 \\
& \quad + 2 a_1 a_2 b_2 - a_1^2 f + 5 a_1 b_0 g - 12 a_2 b_0 l \\
& \quad - 7 a_1 h n + 12 a_2 h m + 2 a_1 l m - 6 g h^2 \\
& \quad + 6 h l^2) \\
& + y_0 y_2 (7 a b_0 c_0 + 2 a b_2 g + 2 a c_1 h - 4 a f l - 7 a m n - 2 a_1^2 c_1 \\
& \quad - 2 a_2^2 b_2 + 4 a_1 a_2 f - 7 a_1 c_0 h - 7 a_2 b_0 g + 17 a_1 g n \\
& \quad + 17 a_2 h m - 10 a_1 l n - 10 a_2 l m - 12 g h l \\
& \quad + 12 l^3) \\
& + y_1^2 \cdot 3 (a b n + a b_0 f + b a_1 g - 2 a b_2 n - 2 b a_2 l \\
& \quad + 2 a_2 b_2 h - a_1 f h - b_0 g h - h^2 n + 2 h l m) \\
& + y_1 y_2 (a b c_0 + 5 a b_0 c_1 - b a_2 g + 3 a b_2 n - 9 a f m \\
& \quad + 2 a_1 b_0 c_0 + 7 a_1 b_2 g - 5 a_1 c_1 h + 4 a_2 b_0 n \\
& \quad - 10 a_2 b_2 l + 11 a_2 f h - 3 c_0 h^2 - 2 a_1 f l \\
& \quad - 3 b_0 g l - 2 a_1 m n + a_2 n^2 - 3 g h m \\
& \quad + 3 h l n + 6 l^2 m)
\end{aligned}$$

$$\begin{aligned}
& + y_2^2 (2ab_0c_0 + 2b_2c_0 - 2ca_1h - 3ac_1m - 2a_2b_0c_0 \\
& - a_2b_2g + 5a_2c_1h + 4a_1c_0m - 2a_1c_1l \\
& + 4a_1fg - 4a_2fl - 3c_0l - 1a_1m^2 + 2a_2m^2 \\
& - 3glmn + 6l^2n) \} \\
& + x_0^3 x_2 \{ y_0^2 \cdot 8 (ac_0h + agm - 2aln - a_1^2c_0 + 2a_1a_2n - a_2^2m \\
& - a_2gh + a_2l^2) \\
& + y_0y_1 (7ab_0c_0 + 2ab_2g + 2ac_1h - 4afl - 7am^2n \\
& - 2a_1^2c_1 - 2a_2^2b_2 + 4a_1a_2f - 7a_1c_0h \\
& - 7a_2b_0g + 17a_1gm + 17a_2hn - 10a_1ln \\
& - 10a_2l^2m - 12a_2l^2 + 12l^3) \\
& + y_0y_2 \cdot 2 (ach - ca_1^2 + 7ac_0m - 2ac_1l + afg - 7an^2 \\
& + 2a_1a_2c_1 - a_2^2f + 5a_2c_0h - 12a_1c_0l \\
& + 12a_1gn - 7a_2gm + 2a_2ln - 6g^2h \\
& + 6g^2n) \\
& + y_1^2 (2abc_0 + ab_0c_1 - 2ba_2g - 3ab_2n - 2a_1b_0c_0 \\
& + 5a_1b_2g - a_1c_1h + 4a_2b_0n - 2a_2b_2l \\
& + 4a_2fh - 4a_1fl - 3b_0gl + 2a_1mn - 4a_2m^2 \\
& - 3l^2m + 6l^2n) \\
& + y_1y_2 (acb_0 + 5ab_2c_0 - ca_1h + 3ac_1m - 9afn
\end{aligned}$$

$$\begin{aligned}
& + 2a_2b_0c_0 - 5a_2b_2g + 7a_2c_1h - 4a_1c_0n \\
& - 10a_1c_1l + 11a_1fg - 3b_0g^2 - 2a_2fl \\
& - 3c_0hl + 4a_1n^2 - 2a_2mn - 3ghn \\
& + 3glm + 6l^2n) \\
& + y_2^2 \cdot 3(acm + ac_0f + ca_2h - 2ac_1n - 2ca_1l + 2a_1c_2g \\
& - a_2fg - c_0gh - g^2n + 2ghn) \} \\
& + x_0^2 x_1^2 \{ y_0^2 \cdot 3(abg - ba_2^2 + 3ab_0n - 2ab_2l + afh - 3am^2 \\
& + 2a_1a_2b_2 - a_1^2f + a_1b_0g - 4a_2b_0l \\
& - 3a_1hn + 4a_2hm + 2a_1lm - 2gh^2 + 2hl^2) \\
& + y_0y_1 \cdot 6(abn + ab_0f + ba_1g - 2ab_2m - 2ba_2l \\
& + 2a_2b_2h + 4a_1b_0n - a_1fh - b_0gh \\
& - 4a_1m^2 - 4b_0l^2 - 5h^2n + 10hlm) \\
& + y_0y_2 \cdot 3(abc_0 + ab_0c_1 - ba_2g - 2b_0c_2n - 2af \\
& + 2a_1c_0 \\
& + 3a_1b_2g - a_1c_1h + 4a_2b_0n - 2a_2b_2l \\
& + 3a_2fh - 3c_0h^2 - 2a_1fl - 7b_0gl \\
& - 2a_1mn - 4a_2m^2 + 5ghm - hln \\
& + 6l^2m) \\
& + f_1^2 \cdot 3(2bf - 2b^2 + 3ba_1n - 2ba_2n + gh - 3b^2 \\
& + 2a_2b_0b_2 + a_1b_0f - b_0^2g - 4a_1b_2m
\end{aligned}$$

$$\begin{aligned}
& -3b_0hn + 4b_2hl + 2b_0lm - 2fh^2 \\
& + 2b_2m^2) \\
& + y_1y_2 \{ 3(abc_1 + ba_1c_0 - ab_2f - ba_2n - bgl + 2a_1b_0c_1 \\
& + 3a_2b_0f - b_0c_0h + 4a_1b_2n - 2a_2b_2n \\
& + 3b_2gl - 3c_1h^2 - 1a_1f - 2b_2gmn \\
& - 2b_0ln - 4b_2l^2 + 5fhl - hmn \\
& + 6l^2m^2) \\
& + y_2^2(abc - ab_2c_1 - ba_2c_0 + 2ca_1b_0 - 3ch^2 + 4a_1b_2c_0 \\
& + 4a_2b_0c_1 - 2a_1c_1n - 6b_2c_0l - 3a_2f - 2 \\
& - 3b_2gl + 3c_0hmn + 3c_1hl + 6fgh \\
& - 3fl^2 - 3gm^2 - 6hn^2 + 12lmn) \} \\
& + x_0^2x_1x_2 \{ y_0^2 \cdot 3(3ab_0c_0 + 2ab_2g + 2ac_1h - 4afl - 3amnv \\
& - 2a_1^2c_1 - 2a_2^2b_2 + 4a_1a_2f - 3a_1c_0h \\
& - 3a_2b_0g + 5a_1gmn + 5a_2hn - 2a_1ln \\
& - 2a_2lm - 4ghl + 4l^3) \\
& + y_0y_1 \cdot 3(abc_0 + 3ab_0c_1 - ba_2g + ab_2n - 5afmn \\
& + 6a_1b_0c_0 + 5a_1b_2g - 3a_1c_1h + 4a_2b_0n \\
& - 6a_2b_2l + 7a_2fh - 7c_0h^2 - 2a_1fl \\
& - 13b_0gl - 6a_1mn - 4a_2m^2 + 9ghn \\
& + hln + 10l^2m)
\end{aligned}$$

$$\begin{aligned}
& + y_0 y_2 \cdot 3 (a c b_0 + 3 a b_2 c_0 - c a_1 h + a c_1 m - 5 a f n \\
& \quad + 6 a_2 b_0 c_0 - 3 a_2 b_2 g + 5 a_2 c_1 h + 4 a_1 c_0 n \\
& \quad - 6 a_1 c_1 l + 7 a_1 f g - 7 b_0 g^2 - 2 a_2 f l \\
& \quad - 13 c_0 h l - 4 a_1 n^2 - 6 a_2 m n + 9 g h n \\
& \quad + g l m + 10 l^2 n) \\
& + f_1^2 \cdot 3 (a b c_1 + 2 b a_1 c_0 - a b_2 f - 3 b g l + 4 a_2 b_0 f \\
& \quad - 2 b_0 c_0 h - 4 a_2 b_2 m + 5 b_2 g h - c_1 h^2 \\
& \quad - 2 a_1 f m - 2 b_0 g m - f h l + 4 l m^2) \\
& + y_1 y_2 (a b c + 8 a b_2 c_1 + 2 b a_2 c_0 + 2 c a_1 b_0 - 9 a f^2 - 3 b g^2 \\
& \quad - 3 c h^2 + 16 a_1 b_2 c_0 + 16 a_2 b_0 c_1 - 12 a_1 c_1 n \\
& \quad - 12 a_2 b_2 n - 6 a_1 f n - 6 a_2 f m \\
& \quad - 18 b_0 g n - 12 b_2 g l - 18 c_0 h m \\
& \quad - 12 c_1 h l + 30 f g h + 6 g m^2 + 6 h n^2 \\
& \quad + 24 l m n) \\
& + y_2^2 \cdot 3 (a c b_2 + 2 c a_2 b_0 - a c_1 f - 3 c h l + 4 a_1 c_0 f \\
& \quad - 2 b_0 c_0 g - 4 a_1 c_1 n - b_2 g^2 + 5 c_1 g h \\
& \quad - 2 a_2 f n - 2 c_0 h n - f g l + 4 l n^2) \}
\end{aligned}$$

This connex may be considered either as a conic or a quartic, depending on whether x or y is the given point. If the curve

has a double point, the curve is singular, but the equation does not. Let

$$z = x_1 = x_2 = 0,$$

so that $(1, 0, 0)$ is a double point, and let this point be the point y . Then

$$\begin{aligned} A^3 x_0^4 y^3 = & x_0^2 x_1^2 \cdot 3(-2gh^2 + 2hl^2) \\ & + x_0^2 x_1 x_2 \cdot 3(-4ghl + 4l^2) \\ & + x_0^2 x_2^2 \cdot 3(-2g^2h + 2gl^2) \\ & + x_0 x_1^3 \cdot 3(-b_0gh - h^2n + 2hlm) \\ & + x_0 x_1^2 x_2 \cdot 3(-c_0h^2 - 2b_0gl - ghm + 4l^2n) \\ & + x_0 x_1 x_2^2 \cdot 3(-b_0g^2 - 2c_0hl - ghn + 4l^2n) \\ & + x_0 x_2^3 \cdot 3(-c_0gh - g^2m + 2gln) \\ & + x_1^4 (bgh - 4bl^2 - b_0^2g - b_0hn + 2b_0lm) \\ & + x_1^3 x_2 (-b_0c_0h + 4b_2gh - 4b_0gm + 2b_0ln - 4b_2l^2 - 3hmn \\ & \quad + 6lm^2) \\ & + x_1^2 x_2^2 (-3b_0gn - 3c_0hm + 6fgh - 6fl^2 - 3gm^2 - 3hn^2 + 12lmn) \\ & + x_1 x_2^3 (-b_0c_0g + 4c_1gh - 4c_0hn + 2c_0lm - 4c_1l^2 - 3gm \\ & \quad + 6ln^2) \\ & + x_2^4 (cgh - 4cl^2 - c_0^2h - c_0gm + 2c_0ln). \end{aligned}$$

Obviously, this curve has a double point at $(1, 0, 0)$ with the same tangents as the quartic. Therefore six of the sixteen intersections of the two curves are used up here. To find the others let us take $(0, 1, 0)$ on (x_1^4) , so that

$$b = 0.$$

Then in order that $(0, 1, 0)$ may also be on $4^3 x_1^4 y^2$ we must have

$$b_0(b_0 g + h n - 2 l m) = 0.$$

If $b_0 = 0$, the intersection is the contact of a tangent from the double point; there are six such. If $b_0 \neq 0$, let us take another intersection which is not one of these contacts as $(0, 0, 1)$. Then

$$c = 0,$$

$$b_0 g + h n - 2 l m = 0,$$

$$c_0 h + g m - 2 l n = 0.$$

These last two equations say that the tangents at the double point, given by

$$h x_1^2 + 2 l x_1 x_2 + g x_2^2 = 0,$$

are copolar to the three points in which x_1 cuts the polar cubic

of the double point, these three points being given by

$$x_0 g_2^3 - 3 \nu g_1 g_2^2 + 3 \nu g_1^2 g_2 - c_1 g_1^3 = 0.$$

But this is true when x_0 is the flex line of the polar curve.

This result can be verified by starting out merely with

$$a = a_1 = a_2 = 0$$

and throwing the polar cubic of $(1, 0, 0)$,

$$3 h x_0^2 x_1^2 + 6 l x_0 x_1 x_2 + 3 g x_0 x_2^2 + l_0 x_1^3 + 3 m x_1^2 x_2 + 3 n x_1 x_2^2 + c_0 x_2^3,$$

into the canonical form by

$$h = g = m = n = 0.$$

Then x_0 is the flex line. It meets $A^3 x^4 y^2$ in points given by

$$-l^2 (l x_1^4 + 4 l_2 x_1^3 x_2 + 6 f x_1^2 x_2^2 + 4 c_1 x_1 x_2^3 + c x_2^4) = 0,$$

evidently, the same points as those in which x_0 meets the

quartic. Therefore the four remaining intersections of $A^3 x^4 y^2$

and $(x_0)^4$ are on the flex line of the polar cubic of the double point.

This result can be verified algebraically. For

$$\begin{aligned} A^3 x^4 y^2 &= (gh - l^2) (6 h x_0^2 x_1^2 + 12 l x_0^2 x_1 x_2 + 6 g x_0^2 x_2^2 + 4 l_0 x_0 x_1^3 + 12 m x_0 x_1^2 x_2 \\ &\quad + 12 n x_0 x_1 x_2^2 + 4 c_0 x_0 x_2^3 + l x_1^4 + 4 l_2 x_1^3 x_2 + 6 f x_1^2 x_2^2 \\ &\quad + 4 c_1 x_1 x_2^3 + c x_2^4) \end{aligned}$$

$$\begin{aligned}
 & - [3hx_0x_1^2 + 6lx_0x_1x_2 + 3gx_0x_2^2 + lx_1^3 + 3mx_1^2x_2 + 3nx_1x_2^2 + ex_2^3] \\
 & \times [4(gh-l^2)x_0 + (l_0g+hn-2lm)x_1 + (e_0h+gm-2ln)x_2] \\
 & = (gh-l^2)(xy)^4 - (xy)^3(dx) \cdot \text{flexline.}
 \end{aligned}$$

The polar curve picks up the six intersections at the double points and above at the six contacts of tangents from the double point, the flex line gives the remaining three.

The flex line can be obtained by a sort of limit process from the polar lines of $(xy)^4$ and $(xy)^3$: instead of at once taking $(1, 0, 0)$ as double point, let us move x_0 the pole line of $(1, 0, 0)$, so that

$$a_1 - a_0 = a.$$

Then if y is $(1, 0, 0)$,

$$\begin{aligned}
 & 3(xy)^2 \cdot (hx)(hy)^5 + (hy)^6 \cdot (dx)(xy)^3 \\
 & = a [4(gh-l^2)x_0 + (l_0g+hn-2lm)x_1 + (e_0h+gm-2ln)x_2].
 \end{aligned}$$

If we divide this by $(xy)^4 = a$ and then let $a=0$ so as to get a double point, we have the flex line.

$A^3 x^4 y^2$ is not the only curve which picks up the six points of contacts of tangents from y when y is a double point. So does $(hx)^5(hy)$ when y is the double point $(1, 0, 0)$

for the coefficient of x_1^5 vanishes when

$$b = b_0 = 0.$$

That it passes through the intersections of $(xy)^4$ and $(xy)^3(xy_1)$ is

$$\begin{aligned} 3(xy)^5(xy_1) &= [5b_0x_1 + 15b_1x_2 + 15b_2x_2^2 + 15b_3x_2^3 + 15b_4x_2^4 + 15b_5x_2^5 \\ &\quad + 4c_0x_0x_2^3 + bx_1^4 + 4b_2x_1^3x_2 + 5fx_1^2x_2^2 + 4c_1x_1x_2^3 + cx_2^4] \\ &\quad \times [4(gh-l^2)x_0 + (b_0g+hn-2lm)x_1 + (c_0h+gm-2ln)x_2] \\ &\quad + [3hx_0x_1^2 + 6lx_0x_1x_2 + 3gx_0x_2^2 + b_0x_1^3 + 3mx_1^2x_2 + 3nx_1x_2^2 + c_0x_2^3] \\ &\quad \times [x_0^2 - 10(gh-l^2) \\ &\quad + x_0x_1(-5b_0g-5hn+10lm) \\ &\quad + x_0x_2(-5c_0h-5gm+10ln) \\ &\quad + x_1^2(-bg-b_0n+2b_2l-fh+n^2) \\ &\quad + x_1x_2(-b_0c_0-2b_2g-2c_1h+4fl+mn) \\ &\quad + x_2^2(-ch-c_0n+2cl-fg+n^2)] \\ &= (xy)^4 \text{ flex line} - (xy)^3(xy_1) \left[\frac{1}{2} A^3 x^2 y^4 + x_0 \cdot \text{flex line} \right], \end{aligned}$$

where by $\frac{1}{2} A^3 x^2 y^4$ is meant the result of forming $A^3 x^2 y^4$ for $y \equiv (1, 0, 0)$ when only $a_1 = a_2 = 0$, dividing by $\frac{1}{2}$, and then letting $a = 0$. Any further pursuit of this relation, however,

to lead to Cayley's identity

$$3(ax)^4(hu)(hy)^5 - (xu)^3(xy) \cdot A^3x^2y^4 + (xu)(xy)^3 \cdot A^3x^4y^2 \\ - 3(xy)^4(hu)^5(hy) \equiv 0.$$

Note 1.

$$H = \sum_{i,j,k} \frac{6!}{i!j!k!} H_{i,j,k} x_0^i x_1^j x_2^k, \quad i+j+k=6,$$

$$H_{600} = agh - al^2 - a_1^2 g - a_2^2 h + 2a, a_2 l,$$

$$H_{510} = \frac{1}{3} (ab_0 g + ahn - 2alm - a_2^2 b_0 - a_1^2 n + 2a, a_2 m \\ - a, gh + a, l^2),$$

$$H_{501} = \frac{1}{3} (ac_0 h + agm - 2aln - a_1^2 c_0 + 2a, a_2 n - a_2^2 m \\ - a_2 gh + a_2 l^2),$$

$$H_{420} = \frac{1}{15} (abg - ba_2^2 + 4ab_0 n - 2ab_2 l + afh - 4am^2 \\ + 2a, a_2 b_2 - a_1^2 f + 2a, b_0 g - 6a_2 b_0 l - 4a, hn \\ + 6a_2 hm + 2a, lm - 3gh^2 + 3hl^2),$$

$$H_{402} = \frac{1}{15} (ach - ca_1^2 + 4ac_0 m - 2ac, l + afg - 4an^2 \\ + 2a, a_2 c, - a_2^2 f + 2a_2 c_0 h - 6a, c_0 l + 6a, gn \\ - 4a_2 gm + 2a_2 ln - 3g^2 h + 3gl^2),$$

$$H_{411} = \frac{1}{15} (2ab_0 c_0 + ab_2 g + ac, h - 2afl - 2amn - a_1^2 c, - a_2^2 b_2 \\ + 2a, a_2 f - 2a, c_0 h - 2a_2 b_0 g + 4a, gm \\ + 4a_2 hn - 2a, ln - 2a_2 lm - 3ghl + 3l^3),$$

$$H_{330} = \frac{1}{10} (abn + ab_0 f + ba, g - 2ab_2 m - 2ba_2 l + 2a_2 b_2 h$$

$$+ 2a_1 b_0 n - a_1 f h - b_0 g h - 2a_1 m^2 - 2b_0 l^2 \\ - 3h^2 n + 6 h l m),$$

$$H_{321} = \frac{1}{30} (abc_0 + 2ab_0 c_1 - b a_2 g - 3afm + 2a_1 b_0 c_0 + 4a_1 b_2 g \\ - 2a_1 c_1 h + 2a_2 b_0 n - 4a_2 b_2 l + 5a_2 f h \\ - 3c_0 h^2 - 2a_1 f l - 6b_0 g l - 2a_1 m n - 2a_2 m^2 \\ + 3ghm + 6l^2 m),$$

$$H_{312} = \frac{1}{30} (ac b_0 + 2ab_2 c_0 - ca_1 h - 3afn + 2a_2 b_0 c_0 - 2a_2 b_2 g \\ + 4a_2 c_1 h + 2a_1 c_0 n - 4a_1 c_1 l + 5a_1 f g \\ - 3b_0 g^2 - 2a_2 f l - 6c_0 h l - 2a_1 n^2 - 2a_2 m n \\ + 3ghn + 6l^2 n),$$

$$H_{222} = \frac{1}{90} (abc + 2ab_2 c_1 + 2b a_2 c_0 + 2c a_1 b_0 - 3af^2 - 3bg^2 - 3ch^2 \\ + 10a_1 b_2 c_0 + 10a_2 b_0 c_1 - 6a_1 c_1 m - 6a_2 b_2 n \\ - 6b_0 c_0 l - 6a_1 f n - 6a_2 f m - 6b_0 g n \\ - 6b_2 g l - 6c_0 h m - 6c_1 h l + 18fgh + 18lmn).$$

The other coefficients may be obtained from these by symmetry.

Note 2.

$$S = \sum \frac{4!}{i!j!k!} S_{ijk} x_0^i x_1^j x_2^k, \quad i+j+k=4,$$

then

$$\begin{aligned} S_{4,0} = & ab_0c_0l - ab_0g^2n - ac_0h^2m + ag^2m^2 + ah^2n^2 - almn \\ & - a_1a_2b_0c_0 + a_1^2c_0m + a_2^2b_0n + a_1b_0g^2 + a_2c_0h^2 \\ & - a_1c_0hl - a_2b_0gl - a_1^2n^2 + a_1a_2mn - a_2^2m^2 \\ & + a_1ghn + a_2ghm - 3a_1ghm - 3a_2hln \\ & + 2a_1l^2n + 2a_2l^2m - g^2h^2 + 2ghl^2 - l^4, \end{aligned}$$

$$\begin{aligned} S_{3,0} = \frac{1}{4} (& abc_0l - abgn - ba_1a_2c_0 - ab_2c_0h + ab_0c_1l + ba_2^2n \\ & - ab_0fg + ba_1g^2 + 2ab_2gm - ac_1h^2m - ba_2gl \\ & - ab_2ln + 2afhn - aflm + a_1^2b_2c_0 - a_1a_2b_0c_1 \\ & + a_2^2b_0f + a_2b_0c_0h + a_1^2c_1m + a_1a_2b_2n - 2a_2^2b_2m \\ & - a_1b_0c_0l + a_2b_2gh + a_2c_1h^2 - 2a_1^2fn + a_1a_2fm \\ & + 2a_1b_0gn - 3a_1b_2gl - a_1c_1hl + 2a_2b_2l^2 - 2a_2b_0l^2 \\ & + a_1fgh - b_0g^2h - 3a_2fhl + 2a_1fl^2 - 2a_1gm^2 \\ & - a_2hmn + b_0gl^2 + a_1lmn + 2a_2lm^2 - gh^2n \\ & + 2ghlm + hl^2n - 2l^3m), \end{aligned}$$

$$S_{3,0,1} = \frac{1}{4} (acl_0l - achm - ca_1a_2b_0 - ab_0c_1g + ab_2c_0l + ca_1^2m$$

$$\begin{aligned}
& -ac_0flv + ca_2lv^2 - ab_2gvn + 2ac_0lvn - ca_1hl \\
& - ac_1lmv + 2afgmv - aflvn - a_1a_2b_2c_0 + a_2^2b_0c_1 \\
& + a_1^2c_0f + a_1b_0c_0g - 2a_1^2c_1v + a_1a_2c_1m + a_2^2b_2v \\
& - a_2b_0c_0l + a_1b_2g^2 + a_1c_1ghv + a_1a_2fvn - 2a_2^2fm \\
& - a_2b_2gl + 2a_2c_0hmv - 3a_2c_1hl - 2a_1c_0lm \\
& + 2a_1c_1l^2 + a_2fflv - c_0gl^2 - 3a_1fgl - a_1f^2v \\
& + 2a_2fl^2 - 2a_2hmv + c_0hl^2 + 2a_1lv^2 + a_2lmv \\
& - g^2hmv + 2ghlv + gl^2v - 2l^3v),
\end{aligned}$$

$$\begin{aligned}
S_{220} = \frac{1}{6} & (-bc_0lv + abc_1l - abf^2 - abv^2 - ab_0b_2c_0 - ba_1a_2c_1 + ab_2^2g \\
& - ab_2c_1v + ba_2^2f - ba_2c_0v + ab_0fvn - ab_2fl \\
& + ba_1gvn - ba_2gmv + ab_2m^2v + ba_2lvn + af^2lv \\
& + bg^2lv - afm^2v - bgl^2v + a_1^2b_2c_1 - a_2^2b_2^2 + a_2b_0^2c_0 \\
& + a_1a_2b_2f + a_1b_2c_0v + a_2b_0b_2g + a_2b_0c_1v \\
& - a_1b_0c_0m - a_1b_0c_1l - a_1^2f^2 - b_0^2g^2 - a_1b_2gmv \\
& - a_2b_0fl + 2a_2b_2hmv + 2a_1b_0v^2 - 3a_1b_2lv \\
& - 3a_2b_0m^2v - a_1fhlv - 2a_2fhlv - b_0ghvn \\
& - 2b_2ghl + 4a_1flm + 4b_0glm - a_1m^2v + 2a_2m^3 \\
& - b_0l^2v + 2b_2l^3 + fghv^2 - fhl^2v - ghvm^2 - lv^2v \\
& + 4hlmv + 3l^2v^2),
\end{aligned}$$

$$\begin{aligned}
S_{2,11} = \frac{1}{12} (& abcl - bca_1a_2 - abe_1g - acb_2h + ba_2^2c_1 + ca_1^2b_2 \\
& - 2ab_0c_0f + ba_1c_0g + ca_2b_0h + 2ab_0c_1n \\
& + 2ab_2c_0m - ba_2c_0l - ca_1b_0l + ab_2fg + ac_1fh \\
& - 2ab_2n^2 - 2ac_1m^2 - af^2l + 2afmn - a_1^2c_1h \\
& - a_2^2b_2f - a_1b_0e_1g - a_2b_2c_0h + 2a_1b_0c_1n \\
& - a_1b_2c_0l + 2a_2b_0c_0m - a_2b_0c_1l + a_1a_2f^2 \\
& + 2a_1c_0fh + 2a_2b_0fg - b_2c_0gh - a_1b_2gm \\
& - 2a_1c_1hn - 2a_2b_2gm - a_2c_1hm - 4a_1c_0 \\
& + 5a_1c_1lm - 4a_2b_0n^2 + 5a_2b_2ln - b_0c_0l^2 \\
& + 2b_2g^2h + 2c_1gh^2 + a_1fgm + a_2fhn - 2b_0 \\
& - 2c_0h^2n - 3a_1fln - 3a_2flm + 4b_0gln \\
& - 2b_2gl^2 + 4c_0hlm - 2c_1hl^2 + 2a_1mn^2 + 2a_2 \\
& - 4fghl - 3ghmn + 4fl^3 + 4glm^2 + 4hln^2 \\
& - 7l^2mn).
\end{aligned}$$

Note 3.

The coefficient of x_0^6 in T is obtained from the expression given by Salmon for the invariant 1 of the cubic as

$$\begin{aligned}
 & a^2 b_0^2 c_0^2 - 6 a^2 b_0 c_0 m n + 4 a^2 b_0 n^3 + 4 a^2 c_0 m^3 - 3 a^2 m^2 n - 6 a a_1 b_0 c_0^2 h \\
 & - 6 a a_2 b_0^2 c_0 g + 6 a a_1 b_0 c_0 g m + 6 a a_2 b_0 c_0 h n + 12 a a_1 b_0 c_0 l n \\
 & + 12 a a_2 b_0 c_0 l m + 4 a b_0^2 g^3 + 4 a c_0^2 h^3 + 12 a b_0 c_0 g h l + 18 a a_1 c_0 h m n \\
 & + 18 a a_2 b_0 g m n - 12 a a_1 b_0 g n^2 - 12 a a_2 c_0 h m^2 - 20 a b_0 c_0 l^3 \\
 & - 24 a a_1 c_0 l m^2 - 24 a a_2 b_0 l n^2 - 12 a b_0 g^2 h n - 12 a c_0 g h^2 m \\
 & - 24 a b_0 g^2 l m - 24 a c_0 h^2 l n + 6 a a_1 g m^2 n - 12 a a_1 h n^3 \\
 & - 12 a a_2 g m^3 + 6 a a_2 h m n^2 + 36 a b_0 g l^2 n + 36 a c_0 h l^2 m \\
 & + 12 a a_1 l m n^2 + 12 a a_2 l m^2 n + 24 a g^2 h m^2 + 24 a g h^2 n^2 \\
 & - 60 a g h l m n + 12 a g l^2 m^2 + 12 a h l^2 n^2 - 12 a l^3 m n \\
 & + 4 a_1^3 b_0 c_0^2 + 4 a_2^3 b_0^2 c_0 - 12 a_1^2 a_2 b_0 c_0 n - 12 a_1 a_2^2 b_0 c_0 m \\
 & + 18 a_1 a_2 b_0 c_0 g h - 3 a_1^2 c_0^2 h^2 - 3 a_2^2 b_0^2 g^2 - 24 a_1^2 b_0 c_0 g l \\
 & - 24 a_2^2 b_0 c_0 h l - 12 a_1^3 c_0 m n + 24 a_1^2 a_2 c_0 m^2 + 24 a_1 a_2^2 b_0 n^2 \\
 & - 12 a_2^3 b_0 m n + 36 a_1 a_2 b_0 c_0 l^2 + 24 a_1^2 b_0 g^2 n + 6 a_1^2 c_0 g h m \\
 & + 6 a_1 a_2 b_0 g^2 m + 6 a_1 a_2 c_0 h^2 n + 6 a_2^2 b_0 g h n + 24 a_2^2 c_0 h^2 m \\
 & + 12 a_1^2 c_0 h l n - 60 a_1 a_2 b_0 g l n - 60 a_1 a_2 c_0 h l m + 12 a_2^2 b_0 g l
 \end{aligned}$$

$$\begin{aligned}
& +12a_1^2c_0l^2m + 12a_2^2b_0l^2n + 8a_1^3n^3 - 12a_1^2a_2mn^2 \\
& - 12a_1a_2^2m^2n + 8a_2^3m^3 - 12a_1b_0g^3h - 12a_2c_0gh^3 + 12a_1c_0gh^2l \\
& + 12a_2b_0g^2hl - 27a_1^2g^2m^2 - 12a_1^2ghn^2 - 6a_1a_2ghmn \\
& - 12a_2^2ghm^2 - 27a_2^2h^2n^2 + 12a_1b_0g^2l^2 + 12a_2c_0h^2l^2 \\
& + 36a_1^2glmn + 36a_1a_2glm^2 + 36a_1a_2hln^2 + 36a_2^2hlmn \\
& - 12a_1c_0hl^3 - 12a_2b_0gl^3 - 24a_1^2l^2n^2 - 12a_1a_2l^2mn - 24a_2^2l^2m^2 \\
& - 12a_1g^2h^2n - 12a_2g^2h^2m + 36a_1g^2hlm + 36a_2gh^2ln \\
& - 12a_1ghl^2m - 12a_2ghl^2n - 36a_1gl^3m - 36a_2hl^3n \\
& + 24a_1l^4n + 24a_2l^4m + 8g^3h^3 - 24g^2h^2l^2 + 24ghl^4 - 8l^6.
\end{aligned}$$

The other coefficients are too difficult to develop in full for the general quartic. They may be obtained in special cases from the coefficient given above (or the symmetrical ones) by means of differential operators.

Note 4

Certain invariants of the quartic have a well-defined geometrical meaning. The two simplest invariants, the A^3 and A^6 , have apolarity meanings and may be included in the list by courtesy. Then comes the A^{15} of Dr. Coble*, which is the condition that the quartic be reducible to the sum of the fourth powers of the six lines of a complete quadrilateral or that the covariant S become two conics; the discriminant, A^{27} ; the A^{48} of Dr. Thomson, expressing that there be a polar conic which is the square of a line; an A^{54} of Dr. Morley's, expressing the condition that the quartic pass thru the vertices of a pentagon; the undulation condition, A^{60} . In this list certain gaps are filled in by the A^{45} , the condition that a polar cubic be made up of three lines, and the A^{51} , the condition that the polar conic of two points be the square of a line.

There should also be an A^{24} , under which the Steineri-

* Ann. Jour. Math., vol. XXXI, p. 357 (1909)

and has the stationary lines of the quartic as double lines.
 For the Steinerian in lines, an $A^{36}f'^8$, and the Cayleyan,
 an $A^{12}f'^8$, touch on the stationary lines.** Then the terms
 of the Steinerian containing only the first powers of $f_i'^4$
 and $(tf)^6$ should be the same as those of the Cayleyan,
 multiplied by an A^{24} to bring them up to the proper
 degree. If this A^{24} vanishes, then Σ has the stationary
 lines as double lines.

**

Proc. Nat. Ac. Sci., Vol. 3, p. 449 (1917).

Vita

Born February 14, 1892, daughter of Benjamin and Rebecca Wiseman Cohen. Graduated from Friends School, Baltimore, Md., in 1909. A.B., Yonkers College, 1912; Resident Fellow, 1912-13. A.M., Johns Hopkins University, 1915; University Scholarship, 1915-16; Fellow, 1917-18.

It is with pleasure that I take this opportunity of expressing my deep gratitude to Professor Morley for his unfailing inspiration and guidance throughout my university career. My thanks and sincere appreciation are also due to Professor Coble and Professor Cohen for their instruction and help, and to Professor Bacon and Professor Lewis of Yonkers College for the splendid foundation they laid for my university studies and for their kindly interest in my work.

Peresa Cohen

